



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁵ : C07H 21/04, C12N 15/63 C12Q 1/68, G01N 27/26	A1	(11) International Publication Number: WO 94/07907 (43) International Publication Date: 14 April 1994 (14.04.94)
(21) International Application Number: PCT/US92/08284 (22) International Filing Date: 25 September 1992 (25.09.92) (71) Applicant: ZOOGEN, INCORPORATED [US/US]; 1105 Kennedy Place, Suite 4, Davis, CA 95616 (US). (72) Inventors: HALVERSON, Joy, Lynn ; DVORAK, Jan ; 2442 Elendil Lane, Davis, CA 95616 (US). (74) Agents: ROWLAND, Bertram, I. et al.; Flehr, Hohbach, Test, Albritton & Herbert, 4 Embarcadero Center, Suite 3400, San Francisco, CA 94111-4187 (US).		(81) Designated States: AU, CA, FI, JP, NO, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, SE). Published <i>With international search report.</i>
(54) Title: AVIAN SEX IDENTIFICATION PROBES (57) Abstract Avian cDNA nucleotide sequences for use as hybridization probes are provided. Probes derived from the sequences find use in defining the sex of a bird, usually by producing hybridization patterns that are sex-specific for most avian species. For a given avian species the probes may hybridize to both Z and W chromosomes so as to differentiate between the two chromosomes on the basis of restriction fragment length polymorphisms. Alternatively, the probes may hybridize exclusively to one of the two sex chromosomes in some species. The hybridization probes taught are also useful for the isolation of other nucleotide sequences that may be used to generate sex-specific hybridization patterns.		

AVIAN SEX IDENTIFICATION PROBES

INTRODUCTION

Technical Field

This invention relates to the field of restriction
5 fragment length polymorphism markers suitable for sex
identification in avian species, and methods for using
such markers.

Background

RFLP (restriction fragment length polymorphism)
10 analysis has proven highly successful in mapping and
discovering new genes in eukaryotic organisms. RFLP
marker probes consist of polynucleotide sequences that
specifically hybridize to a region of the chromosome.
These chromosomal regions of hybridization are revealed
15 to be polymorphic between individuals of the same
species when the chromosomal DNA is digested by
restriction endonucleases and analyzed by hybridization
analysis. Different RFLP alleles are distinguished
from one another on the basis of the hybridization
20 banding patterns produced after size separation.
Genetic linkage analysis between RFLP markers and
uncharacterized genes has proven to be a useful
technique for isolating and mapping uncharacterized
genes of interest. Although the RFLP marker has most
25 frequently been used to identify the chromosomal
disruptions responsible for genetic diseases, it is
also of interest to use RFLP markers in deciphering

other complex genetic regulatory questions, such as sexual development in animals.

While the genetics and biochemistry of sex determination in mammals has been the subject of extensive scientific investigation, similar studies have not been carried out with respect to birds. This is surprising in view of the commercial importance of numerous avian species e.g., chickens and turkeys. It is of interest to provide research tools useful for deciphering the complex process of genetic sex determination in birds. Sex-specific genetic markers are of particular interest. Such markers may be used to sexually identify immature birds prior to the development of gender specific morphological differences. Early sexual identification is an important consideration when breeding those birds that become sexually mature prior to the development of external sexual characteristics. Accordingly, there is interest in providing methods for preventing undesired matings by permitting gender identification (and gender separation) prior to the development of sexual maturity.

Genetic sexual identification is also useful in the breeding of rare bird species with unidentified secondary sexual characteristics; captive breeding programs may thus be effectively organized.

Sex chromosomes, as opposed to autosomal chromosomes, differ with respect to one another in size and genetic composition. Thus, some regions of one sex chromosome contains genes which have no corresponding allele on the other sex chromosome.

One of the principal ways in which the sex chromosomes of birds differ from man and other mammals is that the

female bird is the "heterogametic" sex, having Z and W sex chromosomes. In mammals, the male is the "heterogametic" sex having both X and Y chromosomes whereas the female is "homogametic" having two X chromosomes.

It is of interest to provide RFLP genetic markers suitable for the identification of DNA regions which are diagnostic of the sex of the bird, where the DNA regions may be common to the Z and W chromosomes such that RFLP's unique to each sex chromosomes may be detected; or the DNA region is unique to one of the sex chromosomes, so that the amount or presence of the RFLP will determine the sex of the bird. Such genetic markers permit the identification of the chromosomally specified sex of an individual bird based on analysis of a DNA preparation derived from the bird.

Relevant Literature

Halverson J., Dvorak, J., Flammer, K. 1985. A new method of avian sex determination- identification of the W body by C-banding of erythrocytes. Proceedings of the Annual Meeting of the Association of Avian Veterinarians. 1985. Boulder, Colorado.

Halverson, J., Rauen, K., 1988. The molecular approach to poultry breeding. Proceedings of the Thirty-seventh Western Poultry Disease Conference and Molecular Biology Workshop. 1988. Davis, California.

SUMMARY OF THE INVENTION

Avian nucleotide sequences for use as a source of hybridization probes are provided. Probes derived from these sequences, Tsex (also designated as pMgl) and related sequences, are used in hybridization for sex identification of many avian species. Procedures are given for using the probes so as to sexually identify

individual birds. For a given avian species, the probes may hybridize to both Z and W chromosomes allowing for differentiation between the two chromosomes on the basis of restriction fragment length polymorphisms. Alternatively, the probes may hybridize exclusively to one of the two sex chromosomes in some species, thus permitting gender identification on the basis of sex-specific hybridization intensity. The probes provide a means for identifying the gender of a bird without reliance on morphological characteristics. The probes also find use in recovering and identifying nucleotide sequences homologous to Tsex and sex-specific nucleotide sequences that are genetically linked to genomic Tsex or Tsex-homologous sequences.

DESCRIPTION OF THE SPECIFIC EMBODIMENTS

Novel nucleic acid sequences and methods for using the sequences are provided for determining the chromosomally specified sex of individual birds. The method is based on the ability of hybridization probes derived from the nucleic acid sequences to hybridize with sequences from the sex chromosomes of birds being analyzed for sexual identification. The nucleic acid sequences provided for, namely Tsex, derivatives thereof, and equivalent sequences, are homologous to nucleic acid sequences present on both or one of the Z and W chromosomes of most birds tested. However, there do exist avian species in which Tsex-homologous sequences are only present on one of the two sex chromosomes. For bird DNA, the term homologous when applied to nucleic acid sequences intends nucleic acid sequences capable of hybridizing to each other at a stringency of at least about 25°C below the T_m (T_m is the temperature at which about half the nucleic acid strands are denatured). In referring to "Tsex homologous sequences" it is intended to include the Tsex sequence itself. Most generally used

hybridization protocols may be used with Tsex-derived probes. See Molecular Cloning: A Laboratory Manual, second edition (1989), Sambrook et al., Cold Spring Harbor Laboratories Press, Cold Spring Harbor, N.Y.,
5 for examples of such protocols.

The subject invention permits the identification of the chromosomally specified sex of a given bird based on a hybridization analysis of genomic DNA extracted from the individual bird being gender tested. Suitable
10 birds for chromosomal identification by Tsex hybridization have Tsex-homologous sequences present on one or both sex chromosomes.

The subject invention provides a means for identifying the gender of avian species without relying on
15 morphological sexual characteristics. Thus the subject invention may be used to identify the sex of birds when secondary (external) sexual characteristics are absent or unknown. The identification of sex in the absence of secondary sexual characteristics is useful because
20 in commercial breeding operations it is advantageous to separate sexually mature birds from one another in order to prevent undesired matings. This separation may be difficult to achieve because some avian species become sexually mature prior to the development of
25 obvious external sexual differences. Thus, by providing for a simple and reliable means of gender identification prior to the development of external sexual characteristics in birds, the subject invention finds use in commercial bird breeding. The subject
30 invention finds use in determining the gender of rare bird species in which the gender specific external characteristics are unknown. By permitting the creation of breeding pairs, the subject invention provides a rational means of breeding rare birds.

Hybridization probes derived from Tsex find a variety of uses in addition to their use in sexual identification. Tsex-derived hybridization probes may be used to isolate Tsex-homologous sequences. Such
5 isolation may be achieved by screening recombinant DNA libraries prepared from avian DNA (or cDNA) from species other than turkey (the Tsex sequence being derived from a turkey cDNA). In addition, these Tsex
10 probes may be used to isolate full length cDNAs for Tsex and the Tsex genomic sequences. Similarly, Tsex-derived hybridization probes may also be used in chromosome walking or jumping techniques to isolate coding and noncoding sequences chromosomally proximal, though not necessarily adjoining, to Tsex-homologous
15 sequences. Tsex sequence derived probes may be employed with chromosome walking and jumping techniques as described in a number of commonly available publications, e.g., Sambrook et al., supra.

Additional uses of the subject invention are found in
20 the isolation of sex determination mutations in avian species. Mutations affecting sex determination have found use in manipulating progeny phenotype in the controlled breeding of numerous animals (e.g., linked X chromosomes, autosomal-sex chromosome
25 translocations). The subject invention facilitates the discovery of sex determination mutants by allowing investigators to determine the sexually specified genotype of mutant birds with questionable sexual morphology.

30 The Tsex sequence was obtained from a cDNA library prepared from turkey embryonic poly(A)mRNA. The Tsex sequence is 959 base pairs in length. The sequence of Tsex is displayed in Table 1.

Table 1

```

1  5' AACAGCATCT GATGCTGCAC CCCTTCAGTA TCTGGCTCCC TACTCAGGCT
51  GCTCCATGGG GGAATACTTC AGAGACAATG GGAAACATGC ATTGATCATC
101 TACGATGACT TGTCCAAACA GGCTGTTGCC TACCGTCAGA TGTCTCTGCT
5  151 GCTGCGTCGT CCGCCTGGCC GTGAAGCTTA CCCAGGTGAT GTGTTCTACC
201 TGCACCTCTCG CCTGCTGGAG AGAGCAGCTA AAATGAATGA TTCCTTTGGA
251 GGAGGCTCTC TGA CTGCTTT GCCCGTCATT GAAACTCAGG CTGGTGATGT
301 GTCTGCTTAC ATTCCAACCA ATGTCATCTC CATCACTGAT GGACAGATCT
351 TCTTGGAAC TGA ACTGTTT TACAAAGTA TCCGTCACG CATCAACGTT
10 401 GGTCTGTCTG TGTCCCGTGT GGGTTCTGCT GCTCAGACCA GGGCAATGAA
451 ACAGGTGGCT GGTACCATGA AGCTGGAGCT GGCTCAGTAC CGTGAAGTGG
501 CTGCCTTGC TCAGTTTGGG TCTGATTGG ATGCTGCCAC ACAACAGCTG
551 CTGAATCGTG GTGTGCGTCT GACAGAGCTC CTGAAACAAG GACAGTATGT
601 TCCCATGGCT ATTGAGGAAC AGGTGTCAGT CATCTATCGT GGTGTAAGAG
15 651 GTCAC TTGGA CAAGCTGGAG CCCAGCAAAA TCACTAAATT TGAGAGTGCT
701 TTCCTGGCTC ATGTACTGAG CCAGGACCAG GCCCTCCCTC TCCACCATCA
751 GGACTGAAGG GAAGATCTCT GACCAGACGG AAGCTAAGCT GAAGGAAATA
801 GTCACAAATT TCCTATCTAC TTTTGAGGCA TAAACTCATT ATCTGTTCAA
851 ACAGACCAGG CTGTTTTTGT TGTACGTGC TTTGCCTCCA TCAAAGACCT
20 901 AAACGTATCG AGTGCTTGAA TGTACAGATC TCACTGAGAA TAAAGTTTC
    CATGTAAA3'

```

Sequence displayed from position 1 to end (position 959)

The Tsex sequence may be used for the production of a variety of nucleic acid hybridization probes, also referred to as Tsex-derived probes.

Hybridization probes based on the Tsex sequence find use for many purposes, including sex identification in birds and the isolation of Tsex-homologous sequences from a genetic library. Probes may be either single or double stranded, either RNA or DNA. Probes may be produced by in vitro or in vivo synthesis. Probes may also be produced by a combination of in vitro and in vivo synthesis. Methods of in vitro probe synthesis include organic chemical synthesis processes or enzymatically mediated synthesis, e.g., by means of SP6 RNA polymerase and a plasmid containing the Tsex sequence under the transcriptional control of an SP6 specific promoter. Usually, probes will have a specific complementary sequence of at least 12 nucleotides, more usually at least 14 nucleotides and preferably greater than 50 nucleotides and, more preferably having the entire Tsex sequence.

Probes may bear either complete or partial homology to Tsex. Probes containing partial homology to Tsex will usually have less than 20% mismatch with Tsex, and preferably less than 10% mismatch with Tsex.

- 5 When performing sex determination analysis using Tsex sequence derived probes not actually containing the complete 959 base pair Tsex sequence, the probes are preferably hybridized against DNA preparations from birds of known sex in order to verify that the probes
10 produce the desired sex-specific hybridization pattern. Similarly, when sexing previously untested species of birds with Tsex sequence derived probes, the sex-specific nature of the hybridization patterns produced with Tsex sequence derived probes are preferably
15 verified by hybridization against birds of known sex.

- Probes may be modified by conjugation to a variety of labels which allow for detection of duplex formation between the probe and its complementary target. Labels include radioactive isotopes, ligands, e.g., biotin,
20 enzymes, fluorescers and the like. A wide variety of protocols for labeling probes and detecting duplexes formed between probes and their target hybridization sequences have been described in the literature. See for example, Berger and Kimmel, editors, Guide to
25 Molecular Cloning Techniques (1987) Academic Press Inc., San Diego, CA.

- Probe sequences may be joined to a variety of other nucleic acid sequences. Among these other nucleic acid sequences are vectors such as plasmids, cosmids,
30 phages, and the like. By joining the probe sequence to a vector sequence, probes may be conveniently created, expanded, stored, and modified.

Nucleic acid preparations suitable for hybridization analysis with Tsex-derived probes may be isolated from any portion of a bird's body containing substantially intact nucleic acids. Preferably, nucleic acid preparation sources may be obtained without killing or injuring the bird and may be easily removed from the test bird. Preferred sources of starting material will be feather pulp, blood, or the like.

Tsex-derived hybridization probes may be used to determine the sex of an individual bird based on nucleic acid hybridization analysis. As previously discussed, hybridization analysis demonstrates that Tsex-homologous sequences are present on at least one, and usually both sex chromosomes of numerous avian species. For those species that have Tsex-homologous sequences present on both Z and W sex chromosomes, hereinafter called Tsex-double species, Tsex sequence derived hybridization probes may be used to distinguish male from female DNA preparations on the basis of the presence of at least one additional hybridization band present in the female chromosomal preparations. For those bird species having Tsex-homologous sequences present on only one sex chromosome, hereinafter called Tsex-single species, the genetically specified sex of the bird from which the DNA was prepared may be determined by employing a Tsex-specific probe and semi-quantitatively comparing the degree of hybridization of the probe to a sample with the degree of hybridization of the same probe to a standard of known sex.

Tsex-double species include, but are not limited to: turkeys, chickens, Canada geese, pheasants, zebra finches, lapwings, sandgrouse, murre, caiques (including: white-bellied and black-headed), macaws (including: blue & gold, scarlet, hyacinth, yellow-collar, hahns, greenwing, and military), red-bellied

parrots, amazons (including: Mexican red-headed, orangewing, red lored, mitred, cherry-head, yellow-nape, and blue-front), cockatoos (including: rose-breasted, goffins, umbrella, mollucan, and sulfur-crest), lorries, cockatiels, budgerigars, and rosellas.

Tsex-single bird species include but are not limited to: African grey parrots, Eclectus parrots, Lovebird, and Conures (including: Halfmoon, Patagonian, and Nanday).

- 10 Some bird species are not amenable to sex identification with Tsex-derived probes with the restriction endonucleases employed, but may be capable of identification with an appropriate endonuclease. Tsex-derived probes hybridize to sequences on both the
- 15 W and the Z chromosomes in these species; however, the hybridization patterns produced do not vary between the sexes. The only species tested that are not currently amenable to sex identification with Tsex-derived probes are bald eagles, red tail hawks, ostriches, penguins,
- 20 cinerous vultures, and Russell Griffon vultures.

- The preferred method of sex identification by hybridization analysis with Tsex-derived nucleic acid hybridization probes requires the immobilization of target hybridization sequences onto a solid support.
- 25 Avian sex identification employing the process of nucleic acid hybridization with Tsex-specific probes may utilize solid immobilization supports suitable for hybridization analysis with most generally recognized nucleic acid hybridization procedures. Suitable
- 30 support membranes capable of binding nucleic acids include nitrocellulose, Nytran™, Zetaprobe™ and the like. A variety of protocols for immobilizing nucleic acids to membranes have been described in the literature. See for example, Berger and Kimmel, supra.

The immobilization procedure may be mediated by capillary transfer, electrophoretic transfer, vacuum transfer and the like. The amount of nucleic acid transferred to the solid support membrane will
5 generally be in the range of about .01 μ g to 20 μ g, preferably in the range of about .05-5 μ g, per sample for analysis.

Chromosomal DNA preparations suitable for hybridization with Tsex-derived probes may be digested with
10 restriction endonucleases prior to the transfer of the nucleic acids to an immobilizing support, usually after size separation where the sex is determined based on the difference in size of one or more fragments. Prior to immobilization, chromosomal DNA preparations from
15 Tsex-double species must be digested with at least one restriction endonuclease, whereas chromosomal DNA from Tsex-single species are preferably, though not essentially, digested with at least one restriction endonuclease. Useful restriction endonuclease are
20 selected for their ability to produce a sex-specific hybridization pattern when hybridized against Tsex-derived probes. Where restriction sites are present in the Tsex- or homologous DNA, a number of fragments which bind to the probe may be produced. Digestion
25 with suitable restriction endonucleases produces chromosomal fragments with sizes capable of being separated by electrophoresis.

When DNA preparations for gender determination analysis are subjected to restriction endonuclease digestion,
30 the DNA preparations are size separated by electrophoresis, and transferred in situ from the gel electrophoresis separation medium to a nucleic acid binding support. Electrophoretic separation should proceed to a point where the degree of chromosome
35 digest fragment separation achieved by electrophoresis

is sufficient to separate Tsex-homologous chromosomal fragments from one another to an extent capable of being detected by hybridization analysis. Suitable gel electrophoresis separation media include agarose, 5 polyacrylamide, mixtures thereof, and the like, at a concentration suitable for the separation of nucleic acid fragments to an extent sufficient to reveal size differences between Tsex-homologous DNA fragments. The in situ transfer of nucleic acids to a binding membrane 10 may be achieved by any of the standard procedures for hybridization analysis. Exemplary, but not exclusive of such procedures are Southern blotting and electroblotting.

After immobilization to a solid support, the samples 15 are hybridized against Tsex-derived hybridization probes. Duplexes formed between the labeled probe and the Tsex-homologous sequences in the samples for analysis and any hybridization standards present are subsequently visualized by methods appropriate to the 20 probe label.

When performing chromosomal sex analysis on Tsex-double species, male and female chromosomal DNA hybridization sex standards are optionally, although preferably, present. By hybridization standards, it is intended 25 chromosomal DNA from an individual of known sex and of the same species as the individual undergoing sex determination analysis; the standard is also present in a quantity essentially the same as the DNA sample for analysis.

30 Male and female DNA samples from Tsex-double species may be distinguished from one another by the presence of at least one hybridization band found in the chromosomal preparation of one sex that is not found in preparations of the other sex. Since, frequently one

obtains a plurality of bands of different sizes which bind to the probe, all that is required to distinguish the sex of the subject bird is that there be a band of a size characteristic of a particular sex, where the
5 absence of the band is characteristic of the other sex. Furthermore, the intensity of the hybridization bands in common between male and female chromosomal DNA preparations have greater (usually approximately double) hybridization intensity in male derived samples
10 because of the double dose of target DNA from the two Z chromosomes. Thus, by comparing DNA samples for analysis with each other, or comparing the samples with known male and female standards of the same species, the sex of the individual animal providing the DNA for
15 analysis may be determined.

Sex identification in Tsex-single species is performed essentially the same as in Tsex-double species; however, the restriction endonuclease digestion and electrophoresis separation steps are optional (although
20 preferably performed) rather than required as with the analysis of Tsex-double species. When analyzing Tsex-single species, all DNA samples for analysis and hybridization standards are necessarily present in approximately the same quantity so that the intensity
25 of hybridization produced may be accurately compared among samples and hybridization standards.

In Tsex-single species where Tsex-homologous sequences are only present on the Z chromosome, probe hybridization with chromosomal preparations from males
30 displays greater (approximately double) hybridization intensity than probe hybridization with equal quantities of DNA from female members of the same species.

Tsex-derived probes also find use in isolating other sex-specific sequences. As previously discussed, Tsex-derived probes are useful for sex identification because hybridization with these probes reveals sex-specific hybridization patterns. Since the Tsex probes which hybridize to the W and Z chromosomes are the same sequence, it is apparent that the sex-specific nature of the hybridization must be attributable to chromosomal sequences other than Tsex itself (or Tsex-homologous sequences in species other than turkey). Thus on the chromosome, Tsex-homologous sequences lie close to nucleic acid sequences that differ from one another in a sex-specific manner. These as yet unidentified sex-specific sequences may be detected by employing conventional library screening techniques, including chromosome walking (Bender *et al.*, *J. Mol. Biol.* (1983) 168:17) and chromosome jumping (Poustka *et al.* *Nature* (1987) 325:323), to recover sequences proximal to Tsex-homologous sequences, in which at least the first round of library screening employs a Tsex sequence derived probe. These library screening techniques are discussed at length in Berger and Kimmel, *supra* and Sambrook *et al.*, *supra*.

Genetic libraries for screening with Tsex-derived probes may be prepared in plasmid, cosmid, or YAC (yeast artificial chromosome) vectors or the like. Portions of nucleotide sequences isolated in a first round of screening that are not homologous to Tsex may be labeled and used as probes for a second round of library screening. The process of repeatedly screening a library with newly isolated sequences may be repeated as desired so as to "walk" or "jump" down the chromosome. Progression down the chromosome from the Tsex homologous region may proceed in either direction. The region of the chromosome analyzed by "walking" techniques and the like may extend for a distance

greater than 5000 kb to either side of the Tsex region (or Tsex-homologous region), however preferred distances for walking are in the range of about 100-1000 kb. Chromosomal regions of interest also include
5 introns within the Tsex-homologous genomic sequence. Nucleotide sequences not homologous to Tsex that are isolated during "walking" may be screened for their ability to produce sex-specific hybridization patterns when used as hybridization probes. It is also of
10 interest to engage in chromosome walking for the purpose of isolating sex-specific nucleotide sequences in those bird species that have non-polymorphic Tsex-homologous sequences on both the W and the Z chromosomes, e.g., penguins.

15 For an example of using Tsex to isolate additional sex-specific nucleotide sequences, consider the case of screening a turkey genomic library with a labeled Tsex DNA sequence as a probe. Plasmids from individual clones recovered in the screening may then be
20 restriction mapped and regions of the recovered plasmids not homologous to Tsex isolated. The isolated regions may then be labeled for use as probes in a second round of library screening. Sequences recovered from library screening may also be tested for use as
25 sex-specific probes by labeling the newly isolated sequences and hybridizing the labeled sequences against Southern blots containing restriction digested genomic DNA from both male and female birds (of the same species). Sequences that reveal sex-specific
30 hybridization pattern polymorphisms may find use in sex identification protocols similar to those employing Tsex-derived probes.

Besides using the nucleic acid as an identifier of sex, the expression product may also be used to identify
35 sex. To the extent that the expression products of the

two sequences differ as to epitopes, antisera or monoclonal antibodies may be prepared which will distinguish between the two alleles, when the Tsex or related gene is present on both chromosomes. Where the gene is only on one chromosome, the amount or presence of the protein will indicate the sex of the bird. Various immunoassays may be used to detect the proteins, using radioisotopes, enzymes, chromophores, fluorophores, chemilumescers, or the like, as detectable labels.

Kits may be provided with probes and standards for determining the sex of birds. Thus, by carrying out the assay with the sample and having genomic DNA which may be processed in the same manner, the results may be compared directly. Thus, the kit would comprise, one or more probes, usually one probe, generally labeled, and genomic DNA for one or both sexes of the bird of interest for processing in the assay and comparison with the sample DNA.

The invention now being generally described, the same will be better understood by reference to the following examples which are provided for purposes of illustration only and are not to be considered limiting of the invention unless so specified.

25

EXAMPLES

RNA Isolation

A cDNA library was prepared from poly A mRNA isolated from turkey embryos. All RNA isolation procedures and manipulations were performed in RNase free laboratory articles. In order to remove RNase contamination, glassware was baked at 180° to 190°C overnight and plasticware was treated with a 0.2% diethylpyrocarbonate (DEPC) solution.

8.3 gms of 3, 4, and 5 day old turkey embryos were pooled together and placed in a lysis buffer (7 ml of buffer/1 gm tissue). The lysis buffer is composed of:

4M Guanidinium isothiocyanate

5 .2M Tris pH 7.5

.01M EDTA

5% β mercaptoethanol (w/v)

The tissue was homogenized in a PolytronTM homogenizer at setting "5" for 1 minute. The homogenate was
10 strained through a cheese cloth and a myra cloth filter (a double layer in which the homogenate first passes through the cheese cloth). The filtrate was then centrifuged at 10,000 x g at 25°C for 10 minutes. The supernatant was removed and N lauryl sarcosine was
15 added until a final concentration of 0.5% was reached. Seven volumes of 4 M LiCl were then added. The supernatant was then allowed to sit for 15-20 hours at 4°C. The solution was then centrifuged at 10,000 x g for 90 minutes at 4°C. The supernatant was removed,
20 and the pellet was resuspended in 100 ml of 3 M LiCl. The solution containing the resuspended pellet was then centrifuged at 10,000 x g for 60 minutes at 4°C and the supernatant removed. The pellet was then dissolved in a solubilization buffer by vortexing. RNA
25 solubilization buffer: 0.1% SDS, 0.01 M pH 7.5 Tris, EDTA pH 8 0.5 mM (about 50 ml buffer/10 gm sample). The RNA was then extracted with an equal volume of phenol : chloroform : isoamyl alcohol (25:25:1) solution. The aqueous phase of the extraction was
30 saved and a 3 M potassium acetate solution was added to make a solution with a final concentration of 0.2 molar potassium acetate. 2.2 volumes of 95% ethanol were then added. The solution was dispensed into 30 ml CorexTM tubes. The CorexTM tubes were stored at -20°C
35 overnight. The tubes were then centrifuged at 10,000 x g for 30 minutes at 4°C. The supernatant was removed and the pellet was washed with a 70% ethanol solution.

The tube was allowed to drain and then subsequently dried in a vacuum oven for 20-60 minutes. The pellet was then resuspended in sterile deionized water.

Isolation of Poly A mRNA

5 Poly A mRNA isolated from the total embryonic turkey RNA preparation by oligo(dT)-cellulose was equilibrated with about 2 ml of loading buffer (loading buffer; 20 mM Tris pH 7.5, 0.1 mM EDTA, 0.5 M LiCl, 0.1% SDS). The oligo (dT)-cellulose slurry was poured into a 1.0
10 ml siliconized Pasteur pipette (plugged with siliconized glasswool). The column was washed with 3 volumes each of (1) sterile deionized water, (2) 0.1M NaOH, 5 mM EDTA, (3) sterile deionized water. The column was washed with water until the column effluent
15 had a pH of less than 8.0. The column was then washed with 5 volumes of loading buffer. An aqueous solution containing the total RNA isolated from turkey embryos was heated to 65°C for 5 minutes. Equal volumes of 2x loading buffer was added to the RNA solution and the
20 solution was allowed to cool to room temperature. The solution was then applied to the top of the oligo(dT)cellulose column. The column effluent was collected and heated again to 65° and reapplied to the top of the column. The column was then washed with 5
25 mls of loading buffer. 1 ml aliquots of column effluent were collected. 5 volumes of 0.1 M LiCl buffer (20 mM Tris pH 7.5, 1 mM EDTA, 0.1 M LiCl, 0.1% SDS) were then added to the column. The mRNA fraction was then eluted with 5 volumes of eluting buffer (1mM
30 Tris pH 7.5 1 mM EDTA, 0.05% SDS). 1.0 ml aliquots of column effluent were collected. The column was then rinsed with the deionized water, until the concentration of RNA in the effluent was approximately 0. The A_{260} of each column fraction was determined in
35 order to measure RNA concentration. The fractions were

then stored at -70°C in 70% ethanol and 0.2 M potassium acetate.

cDNA Cloning

5 The embryonic turkey cDNA library was generated using the methods and vectors described by Alexander et al. "Dimer - Primer cDNA Cloning", Methods in Enzymology 154, Academic Press (1987). The cDNA was cloned into the SstI, site of pARC 7. Plasmid pARC 7 was digested with SstI and poly (dT) tailed using terminal
10 deoxynucleotidyl transferase. Isolated embryonic turkey mRNA was allowed to anneal to the poly-dT tails via the poly-A tails of the mRNAs. After annealing to the poly-dT tail primers, the first strand of cDNA synthesis was catalyzed using murine Moloney leukemia
15 virus reverse transcriptase. The first DNA strand was then poly-dG tailed using terminal deoxynucleotidyl transferase. The modified vector bearing cDNA strands on both termini was then digested with the restriction enzyme BamHI. A BamHI linker with a poly-dC tail was
20 then annealed to the poly-dG tail at 42°C. The mixture was then cooled, permitting the constructs to circularize through the annealing of the BamHI sticky ends. Circularization was completed by the use of T4 DNA ligase. The RNA strand still remaining in the
25 construction was removed by mixing the RNA-DNA duplex with RNase H and replacing the RNA with DNA using DNA polymerase I. The remaining nick was then closed with T4 DNA ligase and the constructs were transformed into E. coli strain DH5α.

30 Isolation of pTsex

Clones from the turkey embryonic cDNA library were selected at random and the plasmids within them prepared by a plasmid miniprep procedure (Birnboim and Doly, Nucleic Acids Research (1979) 7:1513). The
35 turkey embryonic cDNA library plasmids were then

screened by digestion with either PstI or XbaI, resulting in the release of the insert portion of the plasmid, and subsequent separation of the vector from the insert by agarose gel electrophoresis. Clones
5 bearing inserts of greater than about 400 base pairs were subjected to further screening.

Further screening was carried out by radioactively labeling the cDNA portion of the isolated clones and using the resulting radioactive probe to screen
10 Southern blots containing restriction digested chromosomal adult turkey DNA from both male and female turkeys. Prior to radioactive labeling, the inserts from the clones for further analysis were subjected to digestion with either PstI or XbaI, followed by
15 separation of the digestion fragments from one another by agarose gel electrophoresis. The plasmid insert was isolated from the gel by electroelution. The insert was then radioactively labeled by nick translation. Nick translation was performed by placing the following
20 mixture in a 1.5 ml eppendorf tube: 0.5 μ g of insert DNA, 2.5 μ l of 10x nick translation buffer (500 mM Tris pH 7.8, 50 mM MgCl₂, 100 mM β mercaptoethanol, 100 μ g/ml BSA), 2.5 μ l 2mM dGTP, 2.5 μ l 2 mM dTTP, 2.5 mM 2.0 mM dCTP, 5 μ l dATP 3000 curies/mM ³²P, 2 μ l DNase I
25 (1U/100 μ l), 1 U DNA polymerase I, H₂O to 25 μ l total volume. The mixture was incubated 75 minutes at 15°C. 8 μ l .25 M EDTA was added, and the mixture was incubated 65°C for 10 minutes to stop the reaction. 1
30 μ l of 10 mg/ml sonicated salmon sperm DNA was then added in order to increase recovery of the probe. The unincorporated nucleotides were removed by gel filtration through a G-75 Sephadex column.

The ³²P labeled inserts were then used as probes to hybridize with a Southern blot containing restriction
35 digested chromosomal DNA prepared from turkey blood.

Southern blots used for the screening of the embryonic turkey cDNA library contained chromosomal DNA from eight turkeys, four males and four females. 5 μ g specimens of chromosomal DNA isolated from the blood of three male and three female turkeys were separately digested with BamHI and EcoR I. Also a single pair of chromosomal DNA preparations from male and female turkeys was digested with HindIII. The chromosomal digest fragments were separated by agarose gel electrophoresis and transferred to a Zetaprobe (BioRad) filter membrane by Southern blotting. All potential gender identification markers were tested against Southern blots containing this combination of restriction digested turkey chromosomal DNA.

The pre-hybridization solution (see section on hybridization conditions for the composition of this solution) was added to a polyethylene baggy containing the Southern blot and allowed to incubate at 65°C overnight. The pre-hybridization solution was removed, and hybridization solution added. The hybridization solution was essentially the same as the pre-hybridization solution with the exception that the hybridization solution contained all of the probe labeled in the nick translation reaction (denatured at 95°C for 10 minutes). The hybridization was allowed to proceed for 36 hours at 65°C.

After hybridization, the membranes were washed by the following procedure. The first wash was in a solution consisting of 1xSSC, 1% SDS, at 45°C for 10 - 15 minutes. The blot was then washed in a solution of 0.1xSSC, 1% SDS, 0.1% sodium pyrophosphate, at 65°C for 40 minutes. After washing, the blot was blotted dry on WhatmanTM 3 MM paper, wrapped in saran wrap, and exposed to KodakTM XAR X-ray film for three days. The X-ray film was subsequently analyzed for differences in

the banding pattern between male and female chromosomal DNA preparations digested with the same restriction enzyme. The desired sex-specific probes should produce no variation in the banding patterns between DNA samples from members of the same sex.

The third embryonic turkey cDNA library clone tested by the above described procedure revealed a sex-specific hybridization pattern. Hybridization with this probe, named the Tsex sequence, with all three male turkey chromosomal DNA preparations digested with BamHI has revealed identical hybridization bands of 19.5 kb and 11.5 kb, whereas hybridization of pTsex with all three BamHI digested chromosomal DNA preparations from female turkeys resulted in the formation of hybridization bands with sizes of 19.5 kb, 13.5 kb, 11.5 kb, 7.3 kb, 6.4 kb and 5.5 kb.

EcoRI digested male chromosomal DNA probed with Tsex resulted in the formation of hybridization bands with sizes of about 13.5 kb, 3.9 kb and 3.4 kb whereas hybridization with EcoRI digested female turkey DNA resulted in the formation of hybridization bands with weights of about 13.5 kb, 9.1 kb, 3.9 kb, 3.4 kb and 2.3 kb. There was no variation in the hybridization patterns produced among the female DNA samples as well as among the male DNA samples.

Male chromosomal DNA digested with HindIII produced hybridization bands with weights of about 6.5 kb and 2.5 kb, whereas hybridization of chromosomal DNA resulted in the formation of hybridization bands with weights of about 6.5 kb, 5.7 kb, 5.3 kb, 4.4 kb, and 2.5 kb.

Blood Collection

Venous blood is collected by clipping a toenail sufficiently short that the blood vessel in the toenail (which usually ends in the distal quarter of the nail) is opened. After the blood has begun to ooze rapidly, a 20 μ l Unopette™ pipette is held to the nail and blood is allowed to flow into the pipette quickly by capillary action. The other end of the Unopette™ fits onto the nozzle of a squirt bottle containing 70% ethanol. The ethanol is squirted through the pipette, washing the collected blood into a 2 ml polypropylene screw cap tube which is then capped. There should be no more than a 30 second delay between blood collection and mixing of the blood with 70% ethanol. The blood is now preserved and will be useful for DNA isolation during several weeks of storage at room temperature.

Blood can also be collected by venipuncture if desired. The wing vein (brachial vein) and jugular vein are preferred. The Unopette™ pipette can be used to transfer 20 μ l of blood into the 2 ml tube for preservation in ethanol as described above. If blood is to be held for any time prior to preservation, it should be immediately chilled in an ice water bath and then held on ice.

Feather Pulp

The proximal shaft of primary feathers that are partially grown contain a mucinous pulp rich in DNA. Feathers are best removed by a firm, steady pulling motion which insures that the feather is removed intact, without leaving a stump. The proximal 2 inches of the feather is cut, and placed in 5 ml of a pre-chilled solution of DNA Isolation Buffer (DIB - 0.1 M NaCl, 0.1mM EDTA, 10 mM Tris pH 7.0). The sample in buffer should be kept on ice until it can be used for DNA preparation.

INTERNATIONAL SEARCH REPORT

International Application No
PCT/GB 96/01341

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES OF USA, vol. 90, no. 18, 15 September 1993, WASHINGTON US, pages 8324-8326, XP002016196</p> <p>✓ 2 R.GRIFFITHS AND B.TIWARI: "The isolation of molecular genetic markers for the identification of sex" cited in the application see the sequence of primer ss2 see page 8324, right-hand column, paragraph 1</p>	3-5, 9-12, 14-18
X	<p>✓ 3 WO,A,94 07907 (ZOOGEN,INCORPORATED) 14 April 1994 see table 2</p>	9-12, 15-18
X	<p>✓ 4 DATABASE WPI Section Ch, Week 8849 Derwent Publications Ltd., London, GB; Class B04, AN 88-348691 XP002016199 & JP,A,63 258 580 (NICHIREI KK) , 26 October 1988 see abstract</p>	9-12, 15-18
X	<p>✓ 5 DATABASE WPI Section Ch, Week 8849 Derwent Publications Ltd., London, GB; Class B04, AN 88-348692 XP002016200 & JP,A,63 258 581 (NICHIREI KK) , 26 October 1988 see abstract</p>	9-12, 15-18
X	<p>✓ 6 PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES OF USA, vol. 90, March 1993, WASHINGTON US, pages 2414-2418, XP002016197 V.DELMAS ET AL.: "A mammalian DNA-binding protein that contains a chromodomain and an SNF2/SWI2-like helicase domain" cited in the application see the whole document</p>	19-21, 24,25
P,X	<p>7 NATURE, vol. 375, 8 June 1995, LONDON GB, page 454 XP002016198 R.GRIFFITHS AND B.TIWARI: "Sex of the last wild Spix's macaw" see the whole document</p>	3-5, 9-12, 14-18

INTERNATIONAL SEARCH REPORT

International Application No

PCT/GB 96/01341

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C12N15/12 C12Q1/68 C07K14/465 C07K14/47 A01K67/027
C07K16/18

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C12Q C07K C12N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	EMBL DATABASE, Accessionnumber D14316 Sequence reference GGJF11 from G. gallus 6 August 1993. XP002016259 compare nucleotides 5-2292 with nucleotides 2221-4508 in figure 5. --- -/--	1-14,19

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

* Special categories of cited documents:

A document defining the general state of the art which is not considered to be of particular relevance

E earlier document but published on or after the international filing date

L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

O document referring to an oral disclosure, use, exhibition or other means

P document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

& document member of the same patent family

Date of the actual completion of the international search

17 October 1996

Date of mailing of the international search report

30. 10. 96

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentaan 2
NL - 2280 HV Rijswijk
Tel. (+ 31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+ 31-70) 340-3016

Authorized officer

Cupido, M

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/GB 96/01341

Patent document: cited in search report	Publication date	Patent family member(s)	Publication date
WO-A-9407907	14-04-94	CA-A- 2124220	14-04-94
		AU-B- 662564	07-09-95
		AU-A- 2696092	26-04-94
		EP-A- 0623139	09-11-94

1/1 - (C) WPI / DERWENT

AN - 88-348691 [49]

AY - 88

XA - C88-154099

XP - N88-264155

DC - B04 C03 D16 P14

FS - CPI

PA - (NICH-) NICHIREI KK

PN - JP63258580 A 881026 DW8849

PR - JP870091976 870416

IC - A01K67/00 ; C07H21/04 ; C12N15/00 ; C12Q1/68

MC - B04-B03B B04-B04A1 B11-C08E B12-K04A6 C04-B03B C04-B04A1 C11-C08E
C12-K04A6 D05-H09 D05-H12

M1 - [01] M423 M710 M781 M903 P831 Q233 V753

UP - 8849

UCC - 8849

AP - JP870091976 870416

XIC - A01K-067/00 ; C07H-021/04 ; C12N-015/00 ; C12Q-001/68

XPN - J63258580

XPR - 87JP-091976

it - CHICKEN DNA DISCRIMINATE HEN COCK LOCALISE FRAGMENT CONTAIN SPECIFIC
NUCLEOTIDE SEQUENCE HEN SPECIFIC SEX CHROMOSOME

AB - J63258580 A DNA fragment containing a specific nucleotide sequence is
localised in hen-specific sex chromosome (W chromosome).S
USE/ADVANTAGE - Discrimination of male and female chickens is done by
professionals. Recent progress of molecular biology can make the
discrimination in molecular level possible. DNA probe is specific for
female chick and thereby accurate discrimination of male and female
chickens is possible. Because the DNA probe recognises hen or game
cock specifically, judgement of unidentified young bird is possible.
(11pp Dwg.No.0/0)

AW - DEOXYRIBONUCLEIC ACID

PAW - (NICH-) NICHIREI KK

TI - Chicken DNA - is used to discriminate between hen and cock, by
localising fragment contg. specific nucleotide sequence in
hen-specific sex chromosome

NC - 001

ORD - 1988-10-26

1/1 - (C) WPI / DERWENT
 AN - 88-348692 [49]
 AY - 88
 XA - C88-154100
 XP - N88-264156
 DC - B04 C03 D16 P14
 FS - CPI
 PA - (NICH-) NICHIREI KK
 PN - JP63258581 A 881026 DW8849
 PR - JP870091977 870416
 IC - A01K67/00 ; C07H21/04 ; C12N15/00 ; C12Q1/68
 MC - B04-B04A1 B11-C08E B12-K04A6 C04-B04A1 C11-C08E C12-K04A6 D05-H01
 D05-H09 D05-H12
 M1 - [01] M423 M760 M903 N102 Q233 V600 V615 V754
 - [02] B615 B701 B713 B720 B815 B831 C811 D011 D012 E720 J011 J171 J521
 L921 M280 M314 M321 M332 M342 M372 M391 M423 M511 M520 M530 M540 M781
 M903 N102 P831 Q233 Q444 Q505 V753; 00945
 - [03] M423 M710 M750 M903 N102 Q233 V753
 M6 - [04] M903 P831 Q233 Q505 R513 R515 R521 R536 R611 R614 R626 R627 R639
 UP - 8849
 UCC - 8849
 AP - JP870091977 870416
 XIC - A01K-067/00 ; C07H-021/04 ; C12N-015/00 ; C12Q-001/68
 XPN - J63258581
 XPR - 87JP-091977
 it - DISCRIMINATE FEMALE MALE TURKEY DNA FRAGMENT CONTAIN SPECIFIC
 NUCLEOTIDE SEQUENCE LOCALISE TURKEY FEMALE SPECIFIC SEX CHROMOSOME
 AB - J63258581 A DNA fragment contains a specific nucleotide sequence
 which is localised in turkey female-specific sex chromosome (W
 chromosome).
 - (1) Turkey female-specific repeat sequence which exists in W
 chromosome emigrates at 0.4kb when electrophoresis is performed after
 digestion with an endonuclease, PstI. (2) Nucleotide sequence of PstI
 0.4kb fragment comprises repeat sequence of about 20 base pairs in
 length. (3) The PstI 0.4kb fragment hybridises female turkey DNA but
 not with male turkey DNA.
 - USE/ADVANTAGE - A DNA probe which is specific for female turkey and
 thereby accurate discrimination of male and female of turkey is
 possible. In addn., because the DNA probe recognises turkey
 specifically, judgement of unidentified young bird, whether it is
 turkey or not, is possible. (7pp Dwg.No.0/0)
 AW - DEOXYRIBONUCLEIC ACID
 PAW - (NICH-) NICHIREI KK
 TI - Discriminating female and male turkey - using DNA fragment contg.
 specific nucleotide sequence localised in turkey female-specific sex
 chromosome
 RRL - 00945
 NC - 001
 ORD - 1988-10-26

ID: GGJF11 , standard; RNA; VRT; 2292 BP.
AC: D14316;
NI: g391639
DT: 06-AUG-1993 (Rel. 36, Created)
DT: 17-FEB-1994 (Rel. 38, Last updated, Version 2)
DE: delta-crystallin enhancer binding protein, complete sequence.
KW: ORF2.
OS: Gallus gallus (chicken)
OC: Eukaryota; Animalia; Metazoa; Chordata; Vertebrata; Aves;
OC: Neornithes; Neognathae; Galliformes; Phasianidae.
RN: [1]
RX: MEDLINE; 94116444.
RA: Funahashi J., Sekido R., Murai K., Kamachi Y., Kondoh H.;
RT: "Delta-crystallin enhancer binding protein deltaEF1 is a zinc
RT: finger-homeodomain protein implicated in postgastrulation

RT: embryogenesis";
RL: Development 119:433-446(1993).
RN: [2]
RP: 1-2292

RA: Funahashi J.;
RT: ;
RL: Unpublished.
CC: Submitted (29-Jan-1993) to DDBJ by: Jun-ichi Funahashi Department
CC: of Molecular Biology School of Science Nagoya University Furo-cho,
CC: Chikusa-ku Nagoya 464-01 Japan Phone: 052-781-5111 x6684 Fax:
CC: 052-782-8575

FH: Key Location/Qualifiers

FH: CDS 257. .1939
FT: /note="beta-gal fusion protein binds to blocks 10 to 3 of
FT: HN fragment of delta-crystallin enhancer."
FT: /product="ORF2"
FT: /codon_start=1

FT: /db_xref="PID:g391640"
FT: /translation="MELKKCCNHCYLIKPPDDNEFYNKQEALQHLIRSSGKLILLDKLL
a.a. 736.
FT: IRLRERGNRVLIFSQMVRMLDILAEYLKYRQFPFQRLDGSIKGELRKQALDHFNAEGSE
FT: DFCFLLSTRAGGLGINLASADTVVIFDSDWNPQNDLQAQARAHRIGQKKQVNIYRLVTK
FT: GSVEEDILERAKKKMVLVDHLVIQRMDDTTGKTVLHTGSTPSSSTPFNKEELSAILKFGAE
FT: ELFKEPEGEEQEPQEMDIDEILKRAETRENEPGPLTVGDELLSQFKVANFSNMDEDDIE
FT: LEPERNSRNWEEIIPESQRRRIEEEEERQKELEEIYMLPRMRNCAKQISFNGSEGRRSRS
FT: RRYSGSDSDSITERKRPKRGRPRPTIPRENIKGFSDAEIRRFIKSYKKFGGPLERLDAV
FT: ARDAELVDKSETDLRRLGELVHNGCIKALKDNSSGQERAGGRLGKVKGPTFRISGVQVN
FT: AKLVISHEEELAPLHKSIPSDPEERKRYVIPCHTKAAHFDIDWGKEDDSNLLVGIYEYA
a.a. 924.
FT: YGSWQCKVNGSRSQNLNTEILPDDPDRTPRQNSYRPVQTTSINY"
FT: source 1. .2292
FT: /organism="Gallus gallus"
FT: /dev_stage="13 day embryo"
FT: /sequenced_mol="cDNA to mRNA"
FT: /tissue_type="lens"
FT: /clone_lib="lambda gt11" . . .

SCORES Init1: 6327 Initn: 8063 Opt: 8924
98.9% identity in 2293 bp overlap

2200 2210 2220 2230 2240 2250
Sa1344 ATTTTCATCATGCCAGAAAAATTTTCCTCCTGGGAAGATTTTGAAGAGGAGCATGGCAAAG

Ggjff11	<div style="text-align: center;"> CCTGGGGAAGATTTTGAAGAGGAGCATGGCAAAG 10 20 30 </div>					
Sa1344	2260	2270	2280	2290	2300	2310
Ggjff11	<div style="text-align: center;"> GAAGAGAGTATGGTTATGCAAGTCTTCACAAAGAGCTTGAACCATTTTTACTAAGAAGAG 40 50 60 70 80 90 </div>					
Sa1344	2320	2330	2340	2350	2360	2370
Ggjff11	<div style="text-align: center;"> TTAAAAAAGATGTAGAAAAGTCTTTACCTGCTAAGGTTGAACAAATTCTGAGGATGGAAA 100 110 120 130 140 150 </div>					
Sa1344	2380	2390	2400	2410	2420	2430
Ggjff11	<div style="text-align: center;"> TGAGTGCATTGCAGAAGCAATATTACAAGTGGATTTTAACAAGGAATTATAAAGCCCTCA 160 170 180 190 200 210 </div>					
Sa1344	2440	2450	2460	2470	2480	2490
Ggjff11	<div style="text-align: center;"> GTAAAGGTTCAAAGGCAGTACCTCAGGCTTCTGAACATTATGATGGAACCTAAGAAGT 220 230 240 250 260 270 </div>					
Sa1344	2500	2510	2520	2530	2540	2550
Ggjff11	<div style="text-align: center;"> GTTGTAACCATTTGCTACCTCATTAAGCCACCAGATGATAATGAATTCTATAATAACAGG 280 290 300 310 320 330 </div>					
Sa1344	2560	2570	2580	2590	2600	2610
Ggjff11	<div style="text-align: center;"> AGGCCTTACAGCATTTGATACGTAGCAGCGGGAACTAATCCTTCTTGACAAGCTACTGA 340 350 360 370 380 390 </div>					
Sa1344	2620	2630	2640	2650	2660	2670
Ggjff11	<div style="text-align: center;"> TTCGTCTGCGAGAACGTGGCAACAGAGTTCTGATTTTCTCTCAGATGGTGAGGATGCTGG 400 410 420 430 440 450 </div>					
Sa1344	2680	2690	2700	2710	2720	2730
Ggjff11	<div style="text-align: center;"> ACATCCTAGCAGAATATCTGAAGTATCGCCAGTTTCCCTTCCAGAGACTTGATGGATCAA 460 470 480 490 500 510 </div>					
Sa1344	2740	2750	2760	2770	2780	2790
Ggjff11	<div style="text-align: center;"> TAAAAGGGGAATTGAGGAAGCAAGCACTGGATCATTTCAATGCAGAAGGATCAGAGGATT 520 530 540 550 560 570 </div>					
Sa1344	2800	2810	2820	2830	2840	2850
Ggjff11	<div style="text-align: center;"> TCTGTTTTTTTACTGTCTACAAGAGCTGGAGGATTAGGTATTAACCTTGGCATCTGCTGACA </div>					

	580	590	600	610	620	630
	2860	2870	2880	2890	2900	2910
Sa1344	CTGTAGTTATTTT	TGATTCTGACTGGAATCCACAGAATGATCTGCAGGCACAGGCGAGAG				
Ggjff11	CTGTAGTTATTTT	TGATTCTGACTGGAATCCACAGAATGATCTGCAGGCACAGGCGAGAG				
	640	650	660	670	680	690
	2920	2930	2940	2950	2960	2970
Sa1344	CTCATAGAATTGGACAGAAGAAACAGGTTAATATTTATCGGCTAGTCACAAAAGGATCAG					
Ggjff11	CTCATAGAATTGGACAGAAGAAACAGGTTAATATTTATCGGCTAGTCACAAAAGGATCAG					
	700	710	720	730	740	750
	2980	2990	3000	3010	3020	3030
Sa1344	TAGAAGAAGATATTCTTGAAAGAGCCAAGAAGAAGATGGTGCTAGACCATTTAGTAATTC					
Ggjff11	TAGAAGAAGATATTCTTGAAAGAGCCAAGAAGAAGATGGTGCTAGACCATTTAGTAATTC					
	760	770	780	790	800	810
	3040	3050	3060	3070	3080	3090
Sa1344	AGAGAATGGACACGACAGGAAAACTGTTCTGCATACAGGTTCAACTCCATCAAGCTCTA					
Ggjff11	AGAGAATGGACACGACAGGAAAACTGTTCTGCATACAGGTTCAACTCCATCAAGCTCTA					
	820	830	840	850	860	870
	3100	3110	3120	3130	3140	3150
Sa1344	CACCTTTTAATAAAGAAGAGTTATCAGCTATTTTGAAGTTTGGTGCTGAGGAACTCTTTA					
Ggjff11	CACCTTTTAATAAAGAAGAGTTATCAGCTATTTTGAAGTTTGGTGCTGAGGAACTCTTTA					
	880	890	900	910	920	930
	3160	3170	3180	3190	3200	3210
Sa1344	AAGAACCTGAAGGAGAAGAACAGGAGCCCCAGGAAATGGATATAGATGAAATCTTGAAGA					
Ggjff11	AAGAACCTGAAGGAGAAGAACAGGAGCCCCAGGAAATGGATATAGATGAAATCTTGAAGA					
	940	950	960	970	980	990
	3220	3230	3240	3250	3260	3270
Sa1344	GAGCTGAAACTCGGGAAAATGAGCCAGGTCCATTGACTGTAGGGGATGAGTTGCTTTCAC					
Ggjff11	GAGCTGAAACTCGGGAAAATGAGCCAGGTCCATTGACTGTAGGGGATGAGTTGCTTTCAC					
	1000	1010	1020	1030	1040	1050
	3280	3290	3300	3310	3320	3330
Sa1344	AGTTCAAGGTGGCGAACTTTTCCAATATGGATGAAGATGATATTGAGTTGGAACCAGAAA					
Ggjff11	AGTTCAAGGTGGCGAACTTTTCCAATATGGATGAAGATGATATTGAGTTGGAACCAGAAA					
	1060	1070	1080	1090	1100	1110
	3340	3350	3360	3370	3380	3390
Sa1344	GAAATTCAAGAAATTGGGAAGAAATCATCCCAGAATCCCAACGGAGAAGGATAGAGGAGG					
Ggjff11	GAAATTCAAGAAATTGGGAAGAAATCATCCCAGAATCCCAACGGAGAAGGATAGAGGAGG					
	1120	1130	1140	1150	1160	1170
	3400	3410	3420	3430	3440	3450
Sa1344	AGGAAAGACAAAAAGAACTTGAAGAAATATACATGCTCCCGAGGATGAGAACTGTGCAA					
Ggjff11	AGGAAAGACAAAAAGAACTTGAAGAAATATACATGCTCCCGAGGATGAGAACTGTGCAA					
	1180	1190	1200	1210	1220	1230
	3460	3470	3480	3490	3500	3510

Sa1344 AACAGATCAGCTTTAATGGGAGTGAAGGAAGACGCAGTAGGAGCAGAAGATATTCTGGAT
 Ggjff11 AACAGATCAGCTTTAATGGGAGTGAAGGAAGACGCAGTAGGAGCAGAAGATATTCTGGAT
 1240 1250 1260 1270 1280 1290
 3520 3530 3540 3550 3560 3570
 Sa1344 CTGATAGTGACTCCATCACAGAAAGAAAACGGCCAAAAAAGCGTGGAAGACCTCGAACCA
 Ggjff11 CTGATAGTGACTCCATCACAGAAAGAAAACGGCCAAAAAAGCGTGGAAGACCTCGAACCA
 1300 1310 1320 1330 1340 1350
 3580 3590 3600 3610 3620 3630
 Sa1344 TTCCTCGAGAAAATATTAAAGGATTTAGTGATGCAGAGATCAGGCGGTTTATCAAGAGTT
 Ggjff11 TTCCTCGAGAAAATATTAAAGGATTTAGTGATGCAGAGATCAGGCGGTTTATCAAGAGTT
 1360 1370 1380 1390 1400 1410
 3640 3650 3660 3670 3680 3690
 Sa1344 ACAAGAAATTTGGTGGCCCTCTGGAAAGGTTAGATGCTGTAGCTAGAGATGCTGAACTGG
 Ggjff11 ACAAGAAATTTGGTGGCCCTCTGGAAAGGTTAGATGCTGTAGCTAGAGATGCTGAACTGG
 1420 1430 1440 1450 1460 1470
 3700 3710 3720 3730 3740 3750
 Sa1344 TTGATAAATCTGAGACAGACCTTAGACGTTTGGGTGAACTTGTAACATAATGGATGCATTA
 Ggjff11 TTGATAAATCTGAGACAGACCTTAGACGTTTGGGTGAGCTTGTAACATAATGGATGCATTA
 1480 1490 1500 1510 1520 1530
 3760 3770 3780 3790 3800 3810
 Sa1344 AGGCTTTAAAGGACAATTCATCTGGACAAGAAAGAGCAGGAGGTAGACTTGGGAAAGTTA
 Ggjff11 AGGCTTTAAAGGACAATTCATCTGGACAAGAAAGAGCAGGAGGTAGACTTGGGAAAGTTA
 1540 1550 1560 1570 1580 1590
 3820 3830 3840 3850 3860 3870
 Sa1344 AAGGCCCAACGTTTCGAATCTCAGGAGTGCAGGTGAATGCAAACTAGTCATCTCTCAG
 Ggjff11 AAGGCCCAACGTTTCGAATCTCAGGAGTGCAGGTGAATGCAAACTAGTCATCTCTCAG
 1600 1610 1620 1630 1640 1650
 3880 3890 3900 3910 3920 3930
 Sa1344 AAGAAGAGCTGGCACCCTGCACAAATCCATTCCTTCAGATCCAGAAGAAAGGAAAAGAT
 Ggjff11 AAGAAGAGCTGGCACCCTGCACAAATCCATTCCTTCAGATCCAGAAGAAAGGAAAAGAT
 1660 1670 1680 1690 1700 1710
 3940 3950 3960 3970 3980 3990
 Sa1344 ATGTCATCCCATGCCACACCAAGGCTGCTCACTTCGATATAGATTGGGGTAAAGAAGATG
 Ggjff11 ACGTCATCCCATGCCACACCAAGGCTGCTCACTTCGATATAGATTGGGGTAAAGAAGATG
 1720 1730 1740 1750 1760 1770
 4000 4010 4020 4030 4040
 Sa1344 ATTCCAATCTGTTAGTAGGCATCTATGAATATGGCTATGGCAGCTGGGAAATGATAA--A
 Ggjff11 ATTCCAATCTGTTAGTAGGCATCTATGAATATGCCTATGGCAGCT-GGCAATGTAAAGTA
 1780 1790 1800 1810 1820 1830
 4050 4060 4070 4080 4090 4100
 Sa1344 AATGGATCCAGATCTCAGCTTAACACAGAAGATTTTACCTGATGATCCAGACAAGAAACC
 Ggjff11 AATGGATCCAGATCTCAGCTTAACACAG-AGATTTTACCTGATGATCCAGAC-AGAATC

1840 1850 1860 1870 1880

4110 4120 4130 4140 4150 4160

Sa1344 CCAGGCAAAGCAGCTACAGACCCGTGCAGACTACCTCATTAATAATTACTGAATAAAGACCT

Ggjf11 CAAGGCAAACAGCTACAGACCCGTGCAGACTACCTCATTAATAATTACTGAATAAAGACCT
1890 1900 1910 1920 1930 1940

4170 4180 4190 4200 4210 4220

Sa1344 TGCAAGAAAGGAAG-CAC-AAAGGCTTGCTGGTGCAGGCAATTCCAAGAGAAGGAAGACA

Ggjf11 TGCAAGAAAGGAAGTCGCTAAAGGCTTGCTGGTGCAGGCAATTCCAAGAGAAGGAAGACA
1950 1960 1970 1980 1990 2000

4230 4240 4250 4260 4270 4280

Sa1344 AGAAATAAGAAGAATAAGATGAAGGCTTCAAAAATAAAAGAAGAAATAAAGAGTGATTCT

Ggjf11 AGGAATAAGAAGAATAAGATGAAGGCTTCAAAAATAAAAGAAGAAATAAAGAGTGATTCT
2010 2020 2030 2040 2050 2060

4290 4300 4310 4320 4330 4340

Sa1344 TCACCACAACCCCTCAGAAAAATCTGATGAAGATGATGAGGAGGAGGATAACAAGGTAAAT

Ggjf11 TCACCACAACCCCTCAGAAAAATCTGATGAAGATGATGAGGAGGAGGATAACAAGGTAAAT
2070 2080 2090 2100 2110 2120

4350 4360 4370 4380 4390 4400

Sa1344 GAAATGAAATCTGAAAATAAAGAAAAATCTAAAAAAATTCCATTGCTGGATACTCCAGTT

Ggjf11 GAAATCAAATCTGAAAATAAAGAAAAATCTAAAAAAATTCCATTGCTGGATACTCCAGTT
2130 2140 2150 2160 2170 2180

4410 4420 4430 4440 4450 4460

Sa1344 CATATTACTGCAACCAGTGAACCAGTTCCTATCTCAGAAGAATCTGAAGAACTCCATCAG

Ggjf11 CATATTACTGCAACCAGTGAACCAGTTCCTATCTCAGAAGAATCTGAAGAACTGGATCAG
2190 2200 2210 2220 2230 2240

4470 4480 4490 4500 4510 4520

Sa1344 AAGACATTTAGTGTGTGCAAAGAAAGAATGAGGCCTGTCAAAGCAGCACTGAAACAGCTG

Ggjf11 AAGACATTTAGTGTGTGCAAAGAAAGAATGAGGCCTGTCAAAG
2250 2260 2270 2280 2290

4530 4540 4550 4560 4570 4580

Sa1344 GATAGACCAGAGAAGGGCCTTTCTGAAAGGGAGCAGCTGGAACATACTAGGCAGTGTCTA